

## Journal Pre-proof

A problem concerning graphs with just three distinct eigenvalues

Peter Rowlinson

PII: S0024-3795(20)30034-3


DOI: <https://doi.org/10.1016/j.laa.2020.01.024>

Reference: LAA 15269

To appear in: *Linear Algebra and its Applications*

Received date: 4 October 2019

Accepted date: 20 January 2020



Please cite this article as: P. Rowlinson, A problem concerning graphs with just three distinct eigenvalues, *Linear Algebra Appl.* (2020), doi: <https://doi.org/10.1016/j.laa.2020.01.024>.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2020 Published by Elsevier.

Rowlinson P (2020) A problem concerning graphs with just three distinct eigenvalues. *Linear Algebra and its Applications*, 592, pp. 260-269. <https://doi.org/10.1016/j.laa.2020.01.024>

© 2020, Elsevier. Licensed under the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International <http://creativecommons.org/licenses/by-nc-nd/4.0/>

A PROBLEM CONCERNING GRAPHS WITH JUST THREE  
DISTINCT EIGENVALUES

Peter Rowlinson<sup>1</sup>  
Mathematics and Statistics Group  
Division of Computing Science and Mathematics  
University of Stirling  
Scotland FK9 4LA  
United Kingdom

**Abstract**

We investigate the problem of finding all the biregular graphs with just three adjacency eigenvalues, one of which is an eigenvalue  $\neq -1, 0$  of maximal possible multiplicity.

*AMS Classification:* 05C50

*Keywords:* biregular graph, eigenvalue multiplicity, main eigenvalue, star complement, strongly regular graph.

---

<sup>1</sup>Tel.: +44 1786 467468; email: p.rowlinson@stirling.ac.uk

# 1 Introduction

We begin with some definitions. Let  $G$  be a graph of order  $n$  with  $(0, 1)$ -adjacency matrix  $A$ . An eigenvalue  $\sigma$  of  $A$  is said to be an eigenvalue of  $G$ , and  $\sigma$  is a *main* eigenvalue if the eigenspace  $\mathcal{E}_A(\sigma)$  is not orthogonal to the all-1 vector in  $\mathbb{R}^n$ . Always the largest eigenvalue, or *index*, of  $G$  is a main eigenvalue, and it is the only main eigenvalue if and only if  $G$  is regular. We say that  $G$  is *integral* if every eigenvalue of  $G$  is an integer. If  $G$  is connected with an eigenvalue  $\neq -1, 0$  of multiplicity  $k$  and comultiplicity  $t = n - k > 1$  then  $k \leq \frac{1}{2}t(t-1)$  [1, Theorem 2.3]. If  $k = \frac{1}{2}t(t-1)$  then we say that  $G$  has an eigenvalue  $\neq -1, 0$  of *maximal multiplicity*.

A *biregular* graph is a graph with precisely two distinct degrees. A *strongly regular* graph, with parameters  $n, r, e, f$ , is an  $r$ -regular graph of order  $n$  in which any two adjacent vertices have  $e$  common neighbours, and any two non-adjacent vertices have  $f$  common neighbours. We take strongly regular graphs to include cliques and co-cliques. In accordance with [9] we say that  $G$  has a *strongly regular decomposition* if its vertex set has a bipartition  $V(G) = X_1 \dot{\cup} X_2$  such that the subgraphs induced by  $X_1$  and  $X_2$  are strongly regular.

Let  $\mathcal{B}$  be the class of connected biregular graphs with just three distinct eigenvalues, and let  $\mathcal{C}$  be the class of connected graphs with an eigenvalue  $\mu \neq -1, 0$  of maximal multiplicity. It is an open problem to determine the graphs in  $\mathcal{B}$  [3], and another open problem, with origins in [1], to determine the graphs in  $\mathcal{C}$ . Graphs with just three distinct eigenvalues, subject to various restrictions, have recently been investigated in [3, 4, 12, 13].

It was noted in [12] that  $\mathcal{B} = \mathcal{C}_1 \cap \mathcal{C}_2$ , where  $\mathcal{C}_1$  is the class of connected graphs with just three distinct eigenvalues, and  $\mathcal{C}_2$  is the class of connected graphs with exactly two main eigenvalues; moreover any graph  $G$  in  $\mathcal{B}$  is either integral or complete bipartite. Many examples are given in [3, 12, 13]. Since the spectrum of a bipartite graph is symmetric about 0, the only bipartite graph in  $\mathcal{B} \cap \mathcal{C}$  is  $K_{1,2}$ . Thus any other graph in  $\mathcal{B} \cap \mathcal{C}$  is non-bipartite and integral.

Apart from  $K_{1,2}$ , the only known graph in  $\mathcal{C}$  is the graph  $\Gamma_{36}$  (of order 36) obtained from  $L(K_9)$  by switching with respect to  $K_8$ ; this is the largest exceptional graph, denoted by  $G473$  in [6]. Since  $\Gamma_{36}$  has spectrum  $-2^{(28)}, 5^{(7)}, 21$ , the only known graphs in  $\mathcal{B} \cap \mathcal{C}$  are  $K_{1,2}$  and  $\Gamma_{36}$ . The problem investigated in this paper is to determine whether there are any other graphs in  $\mathcal{B} \cap \mathcal{C}$ . A solution to the problem remains elusive, but we do obtain two restrictive conditions on the graphs in question: in Proposition 4.2 we provide a relation between degrees and spectrum, and in Theorem 4.4 we show that such graphs have a strongly regular decomposition.

We use the notation of the monograph [7], where the basic properties of graph spectra can be found in Chapter 1. We write  $\mathbf{j}$  for an all-1 vector, its length determined by context; and we write  $u \sim v$  to indicate that vertices  $u, v$  of a graph are adjacent.

## 2 Graphs in $\mathcal{B}$

Here we note the results we require from [3] and [12]. Let  $G$  be a graph in  $\mathcal{B}$  with degrees  $d_1, d_2$ , and let  $X_i = \{v \in V(G) : \deg(v) = d_i\}$ ,  $|X_i| = n_i$  ( $i = 1, 2$ ). Let the distinct eigenvalues of  $G$  be  $\rho, \lambda, \mu$ , where  $\rho$  is the index of  $G$ . We have

$$(A - \lambda I)(A - \mu I) = \mathbf{v}\mathbf{v}^\top, \quad (1)$$

where  $A\mathbf{v} = \rho\mathbf{v}$ . Let  $\mathbf{v} = (v_1, v_2, \dots, v_n)^\top$ : since  $G$  is connected, we may take  $v_i > 0$  ( $i = 1, 2, \dots, n$ ). It follows from Eq.(1) that  $v_i^2 = d_1 + \lambda\mu$  if  $i \in X_1$  and  $v_i^2 = d_2 + \lambda\mu$  if  $i \in X_2$ . Hence  $\mathbf{v} = \begin{pmatrix} a_1 \mathbf{j} \\ a_2 \mathbf{j} \end{pmatrix}$ , where  $a_i^2 = d_i + \lambda\mu$  ( $i = 1, 2$ ) and the partition of  $\mathbf{v}$  is determined by  $X_1 \dot{\cup} X_2$ .

**Lemma 2.1** *If  $G \in \mathcal{B}$  then either*

- (a)  $\mu$  is non-main and  $a_1 a_2 = -\lambda(\mu + 1)$ , or
- (b)  $\lambda$  is non-main and  $a_1 a_2 = -\mu(\lambda + 1)$ .

**Proof.** See [12, Lemma 2.2].

**Lemma 2.2** *If  $G \in \mathcal{B}$  then the sets  $X_1, X_2$  form an equitable bipartition of  $G$  with divisor matrix  $K = \begin{pmatrix} k_{11} & k_{12} \\ k_{21} & k_{22} \end{pmatrix}$ , where*

$$k_{11} = \frac{a_1 \rho - a_2 d_1}{a_1 - a_2}, \quad k_{12} = a_1 \frac{d_1 - \rho}{a_1 - a_2}, \quad k_{21} = a_2 \frac{d_2 - \rho}{a_2 - a_1}, \quad k_{22} = \frac{a_2 \rho - a_1 d_2}{a_2 - a_1},$$

$$|X_1| = \frac{(\rho - a_2^2 + \lambda\mu)(\rho - \lambda)(\rho - \mu)}{(\rho + \lambda\mu + a_1 a_2)(a_1^2 - a_1 a_2)}, \quad |X_2| = \frac{(\rho - a_1^2 + \lambda\mu)(\rho - \lambda)(\rho - \mu)}{(\rho + \lambda\mu + a_1 a_2)(a_2^2 - a_1 a_2)}.$$

**Proof.** See [3, Theorem 4.3].

## 3 Graphs in $\mathcal{C}$

In this section we consider a graph  $G$  of order  $n = \frac{1}{2}t(t+1)$  with an eigenvalue  $\mu \neq -1, 0$  of multiplicity  $k = \frac{1}{2}t(t-1)$  and comultiplicity  $t > 1$ . Let  $X$  be a star set for  $\mu$  in  $G$ , that is, a set of  $k$  vertices such that  $\mu$  is not an eigenvalue of  $G - X$ . The induced subgraph  $H = G - X$  is called the corresponding star complement for  $\mu$  (see [7, Chapter 5]). By [7, Proposition 5.1.4] the  $H$ -neighbourhoods of vertices in  $X$  are non-empty and distinct. If  $A_X$  is the adjacency matrix of the subgraph induced by  $X$  then  $G$  has adjacency matrix  $A = \begin{pmatrix} A_X & B^\top \\ B & C \end{pmatrix}$ , where  $C$  is the adjacency matrix of  $H$ . Then

$$\mu I - A_X = B^\top (\mu I - C)^{-1} B \quad (2)$$

by [7, Theorem 5.1.7]. It follows that if  $S = (B|C - \mu I) = (\mathbf{s}_1|\mathbf{s}_2|\dots|\mathbf{s}_n)$  then  $\mu I - A = S^\top(\mu I - C)^{-1}S$ , and if  $\langle\langle \mathbf{x}, \mathbf{y} \rangle\rangle$  denotes the bilinear form  $\mathbf{x}^\top(\mu I - C)^{-1}\mathbf{y}$  ( $\mathbf{x}, \mathbf{y} \in \mathbb{R}^t$ ) then

$$\langle\langle \mathbf{s}_u, \mathbf{s}_v \rangle\rangle = \begin{cases} -1 & \text{if } u \sim v \\ 0 & \text{if } u \not\sim v \text{ and } u \neq v \\ \mu & \text{if } u = v. \end{cases}$$

From [1] we know that the functions  $\langle\langle \mathbf{s}_1, \mathbf{x} \rangle\rangle^2, \dots, \langle\langle \mathbf{s}_n, \mathbf{x} \rangle\rangle^2$  form a basis for the space of homogeneous quadratic functions on  $\mathbb{R}^t$ . In particular, there exist  $\alpha_1, \dots, \alpha_n \in \mathbb{R}$  such that

$$\langle\langle \mathbf{x}, \mathbf{x} \rangle\rangle = \sum_{u=1}^n \alpha_u \langle\langle \mathbf{s}_u, \mathbf{x} \rangle\rangle^2. \quad (3)$$

Taking  $\mathbf{x} = \mathbf{s}_v$  in Eq.(3) we have  $\mu = \alpha_v \mu^2 + \sum_{u \sim v} \alpha_u$ . Thus if  $\mathbf{a} = (\alpha_1, \dots, \alpha_n)^\top$  then

$$\mu \mathbf{j} = (\mu^2 I + A) \mathbf{a}. \quad (4)$$

It also follows from Eq.(3) that  $\langle\langle \mathbf{x}, \mathbf{y} \rangle\rangle = \sum_{u=1}^n \alpha_u \langle\langle \mathbf{s}_u, \mathbf{x} \rangle\rangle \langle\langle \mathbf{s}_u, \mathbf{y} \rangle\rangle$ ; in particular, taking  $\mathbf{y} = \mathbf{s}_v$  we have  $\langle\langle \mathbf{s}_v, \mathbf{x} \rangle\rangle = \alpha_v \mu \langle\langle \mathbf{s}_v, \mathbf{x} \rangle\rangle - \sum_{u \sim v} \alpha_u$ . Hence if  $D = \text{diag}(\alpha_1, \dots, \alpha_n)$  then

$$\begin{pmatrix} \langle\langle \mathbf{s}_1, \mathbf{x} \rangle\rangle \\ \langle\langle \mathbf{s}_2, \mathbf{x} \rangle\rangle \\ \vdots \\ \langle\langle \mathbf{s}_n, \mathbf{x} \rangle\rangle \end{pmatrix} = (\mu I - A) D \begin{pmatrix} \langle\langle \mathbf{s}_1, \mathbf{x} \rangle\rangle \\ \langle\langle \mathbf{s}_2, \mathbf{x} \rangle\rangle \\ \vdots \\ \langle\langle \mathbf{s}_n, \mathbf{x} \rangle\rangle \end{pmatrix}.$$

Taking  $\mathbf{x} = \mathbf{s}_1, \dots, \mathbf{s}_n$ , we obtain:

**Proposition 3.1** *If  $G \in \mathcal{C}$  then, with the notation above,*

$$\mu I - A = (\mu I - A) D (\mu I - A). \quad (5)$$

## 4 Graphs in $\mathcal{B} \cap \mathcal{C}$

Here we retain the notation of Sections 2 and 3, and take  $G$  to be a graph in  $\mathcal{B} \cap \mathcal{C}$  other than  $K_{1,2}$ . Let  $\mu$  be the eigenvalue  $\neq -1, 0$  of maximal multiplicity  $\frac{1}{2}t(t-1)$ . Note that  $t > 2$  because  $G$  has order  $> 3$ . By [1, Theorem 3.1],  $\mu$  is a main eigenvalue, and so the main eigenvalues of  $G$  are  $\rho$  and  $\mu$ . It follows from Lemma 2.1 that  $a_1 a_2 = -\mu(\lambda + 1)$ , and it follows from [11, Proposition 2.1] that

$$(A - \rho I)(A - \mu I) \mathbf{j} = \mathbf{0}. \quad (6)$$

**Lemma 4.1** *The matrix  $\mu^2 I + A$  is invertible.*

**Proof.** Suppose by way of contradiction that  $\lambda = -\mu^2$ . Note first that if the eigenvalues  $\mu_1, \dots, \mu_t$  of  $G$  other than  $\mu$  have mean  $d$  then  $k\mu + td = 0$  and  $k\mu^2 + \sum_{i=1}^t \mu_i^2 = n\bar{d}$ , where  $\bar{d}$  is the mean degree in  $G$ . We have

$$\sum_{i=1}^t (\mu_i - d)^2 = \sum_{i=1}^t \mu_i^2 - td^2 = (t+k)\bar{d} - \mu^2(k + t(\frac{k^2}{t^2})) = (t+k)\bar{d} - \mu^2 \frac{k}{t}(t+k),$$

and so  $k\mu^2/t \leq \bar{d}$ . Since  $k = \frac{1}{2}t(t-1)$  and  $\bar{d} < \rho$  we have  $(t-1)\mu^2 < 2\rho$ . Now  $\rho = (t-1)\mu^2 - \frac{1}{2}t(t-1)\mu$ , whence  $\mu > 0$ , for otherwise  $-\mu \geq 2$  and we find that  $\rho > 2n$ . Since  $\rho = \frac{1}{2}(t-1)(2\mu-t)\mu$  we have  $\mu > \frac{1}{2}t$ . Therefore  $\frac{1}{4}t^2(t-1) < (t-1)\mu^2 < 2\rho \leq 2n-2 = t(t+1)-2$ , whence  $t \leq 5$  and  $n \leq 15$ . Now we have a contradiction either from [3, Table 1] or by calculating  $\rho, \mu, n$  when  $t = 5, 4, 3$ .  $\square$

In view of Lemma 4.1, we have  $\mathbf{a} = \mu(\mu^2 I + A)^{-1} \mathbf{j}$  from Eq.(4). Now  $\mu(\mu^2 I + A)^{-1}$  is a polynomial in  $A$ , while Eq.(6) shows that  $\mathbf{j}$  is annihilated by a quadratic in  $A$ . It follows that  $\mathbf{a} \in W$ , where  $W = \langle \mathbf{j}, A\mathbf{j} \rangle$ , i.e.  $W$  is the subspace spanned by  $\begin{pmatrix} \mathbf{j} \\ \mathbf{j} \end{pmatrix}$  and  $\begin{pmatrix} d_1 \mathbf{j} \\ d_2 \mathbf{j} \end{pmatrix}$ . Hence there exist  $\alpha, \beta \in \mathbb{R}$  such that  $\alpha_u = \alpha + \beta d_1$  if  $u \in X_1$  and  $\alpha_u = \alpha + \beta d_2$  if  $u \in X_2$ . Writing  $\mathbf{d} = A\mathbf{j}$ , we have from Eq.(4) and Eq.(6):

$$\begin{aligned} \mu \mathbf{j} &= \mu^2 \mathbf{a} + A\mathbf{a} = \mu^2 \mathbf{a} + \alpha A\mathbf{j} + \beta A\mathbf{d} = \mu^2(\alpha \mathbf{j} + \beta \mathbf{d}) + \alpha \mathbf{d} + \beta A^2 \mathbf{j} \\ &= \mu^2(\alpha \mathbf{j} + \beta \mathbf{d}) + \alpha \mathbf{d} + \beta((\rho + \mu)\mathbf{d} - \rho \mu \mathbf{j}) = (\mu^2 \alpha - \beta \rho \mu) \mathbf{j} + (\alpha + \mu^2 \beta + \beta \mu + \beta \rho) \mathbf{d}. \end{aligned}$$

It follows that

$$\alpha = \frac{\mu^2 + \mu + \rho}{(\mu + 1)(\mu^2 + \rho)}, \quad \beta = \frac{-1}{(\mu + 1)(\mu^2 + \rho)}. \quad (7)$$

**Proposition 4.2** *If  $G \in \mathcal{B} \cap \mathcal{C}$  then  $(\rho - d_1)(d_2 - \rho) = (\mu + 1)^2 \rho$ .*

**Proof.** Note that  $W (= \langle \mathbf{j}, A\mathbf{j} \rangle)$  is  $D$ -invariant and  $A$ -invariant. We find the matrix of  $D(\mu I - A)|_W$  with respect to the basis  $\{\mathbf{j}, \mathbf{d}\}$ . First,  $D(\mu I - A)\mathbf{j} = D \begin{pmatrix} (\mu - d_1)\mathbf{j} \\ (\mu - d_2)\mathbf{j} \end{pmatrix} = \begin{pmatrix} w_1 \mathbf{j} \\ w_2 \mathbf{j} \end{pmatrix}$ , where  $w_i = (\alpha + \beta d_i)(\mu - d_i)$  ( $i = 1, 2$ ). Now  $\begin{pmatrix} w_1 \mathbf{j} \\ w_2 \mathbf{j} \end{pmatrix} = p \begin{pmatrix} \mathbf{j} \\ \mathbf{j} \end{pmatrix} + q \begin{pmatrix} d_1 \mathbf{j} \\ d_2 \mathbf{j} \end{pmatrix}$ , where  $p = \frac{w_2 d_1 - w_1 d_2}{d_1 - d_2}$  and  $q = \frac{w_1 - w_2}{d_1 - d_2}$ . Secondly,  $D(\mu I - A)\mathbf{d} = D(\rho \mu I - \rho A)\mathbf{j}$  by Eq.(6), and so the required matrix is  $\begin{pmatrix} p & \rho p \\ q & \rho q \end{pmatrix}$ .

Now Eq.(5) shows that  $D(\mu I - A)|_W$  is idempotent and so  $p + \rho q = 1$ , i.e.  $d_1 - d_2 = (\rho - d_2)(\alpha + \beta d_1)(\mu - d_1) - (\rho - d_1)(\alpha + \beta d_2)(\mu - d_2)$ , or  $1 = \beta(\rho - d_1)(\rho - d_2) - (\rho - \mu)(\rho\beta + \alpha)$ . Substituting for  $\alpha, \beta$  from Eq.(7), we obtain

$$\frac{(\rho - d_1)(\rho - d_2)}{(\mu + 1)(\mu^2 + \rho)} + \frac{(\rho - \mu)\mu}{\mu^2 + \rho} = -1,$$

and the result follows.  $\square$

**Example 4.3.** For the graph  $\Gamma_{36}$  (with  $d_1 < d_2$ ) we have  $\mu = -2$ ,  $\lambda = 5$ ,  $\rho = 21$ ,  $d_1 = 18$ ,  $d_2 = 28$ ,  $a_1 = 2\sqrt{2}$ ,  $a_2 = 3\sqrt{2}$ ,  $\alpha = -23/25$ ,  $\beta = 1/25$ .

For  $i = 1, 2$ , let  $G_i$  be the subgraph of  $G$  induced by  $X_i$ , with adjacency matrix  $A_i$ . In what follows, the neighbourhood of a vertex  $v$  in  $G$  is denoted by  $\Delta(v)$ . We also write  $A = (a_{ij})$  and  $A^2 = (a_{ij}^{(2)})$ .

**Theorem 4.4** *If  $G \in \mathcal{B} \cap \mathcal{C}$  then the graph  $G_1$  is strongly regular with parameters  $n_1, k_{11}, e_1, f_1$ , where*

$$(d_1 - d_2)e_1 = (\mu^2 + \mu + \rho - d_2)\lambda(\mu + 1) + d_1\mu^2 + 3d_1\mu + d_1\rho - d_1d_2 - d_2\mu + \rho,$$

$$(d_1 - d_2)f_1 = (\mu^2 + \mu + \rho - d_2)(d_1 + \lambda\mu);$$

*and the graph  $G_2$  is strongly regular with parameters  $n_2, k_{22}, e_2, f_2$ , where*

$$(d_2 - d_1)e_2 = (\mu^2 + \mu + \rho - d_1)\lambda(\mu + 1) + d_2\mu^2 + 3d_2\mu + d_2\rho - d_1d_2 - d_1\mu + \rho,$$

$$(d_2 - d_1)f_2 = (\mu^2 + \mu + \rho - d_1)(d_2 + \lambda\mu).$$

**Proof.** From Eq.(3) we have

$$\langle \mathbf{x}, \mathbf{y} \rangle = (\alpha + \beta d_1) \sum_{u \in X_1} \langle \mathbf{s}_u, \mathbf{x} \rangle \langle \mathbf{s}_u, \mathbf{y} \rangle + (\alpha + \beta d_2) \sum_{u \in X_2} \langle \mathbf{s}_u, \mathbf{x} \rangle \langle \mathbf{s}_u, \mathbf{y} \rangle,$$

where  $\alpha, \beta$  are given by Eq.(7). Hence if  $i, j \in X_1$  and  $i \sim j$  then

$$-1 = -2(\alpha + \beta d_1)\mu + (\alpha + \beta d_1)|\Delta(i) \cap \Delta(j) \cap X_1| + (\alpha + \beta d_2)|\Delta(i) \cap \Delta(j) \cap X_2|. \quad (8)$$

On the other hand, since  $a_{ij}^{(2)} = a_1^2 + \lambda + \mu$  we have

$$d_1 + \lambda + \mu + \lambda\mu = |\Delta(i) \cap \Delta(j) \cap X_1| + |\Delta(i) \cap \Delta(j) \cap X_2|. \quad (9)$$

Now we can solve (8) and (9) for  $|\Delta(i) \cap \Delta(j) \cap X_1|$  and  $|\Delta(i) \cap \Delta(j) \cap X_2|$ , to find that the number of common neighbours of  $i$  and  $j$  in  $X_1$  is  $e_1$ , where  $(d_1 - d_2)e_1$  is as given in the statement of the Proposition. Again if  $i, j \in X_1$  and  $i \not\sim j$  then

$$0 = (\alpha + \beta d_1)|\Delta(i) \cap \Delta(j) \cap X_1| + (\alpha + \beta d_2)|\Delta(i) \cap \Delta(j) \cap X_2|. \quad (10)$$

and

$$d_1 + \lambda\mu = |\Delta(i) \cap \Delta(j) \cap X_1| + |\Delta(i) \cap \Delta(j) \cap X_2|. \quad (11)$$

From (10) and (11), we find that the number of common neighbours of  $i$  and  $j$  in  $X_1$  is  $f_1$ , where  $(d_1 - d_2)f_1$  is as stated above. We may interchange  $X_1$  and  $X_2$  to obtain the second part of the Proposition.  $\square$

Note that Theorem 4.4 says that  $G$  has a strongly regular decomposition in the sense of [9]. For  $\Gamma_{36}$  (with  $d_1 < d_2$ ) we have  $G_1 = L(K_8)$  with  $e_1 = 6$ ,  $f_1 = 4$ ,  $k_{11} = 12$ , and  $G_2 = K_8$  with  $e_2 = 6$ ,  $f_2 = 9$ ,  $k_{22} = 7$ . (Although  $f_2 > k_{22}$ , this is not a contradiction because  $G_2$  has no non-adjacent vertices, and  $(A - 7I)(A + I)(A + 2I) = O$ .)

**Proposition 4.5** *Let  $G$  be a graph in  $\mathcal{B} \cap \mathcal{C}$  other than  $K_{1,2}$ . If  $\mu$  is not an eigenvalue of both  $G_1$  and  $G_2$  then  $G = \Gamma_{36}$ .*

**Proof.** Any non-main eigenvalue of  $G_i$  is a root of  $x^2 - (e_i - f_i)x - (k_{ii} - f_i)$  (cf. [7, Eq.(3.17)]), and one can check that  $\mu^2 - (e_i - f_i)\mu - (k_{ii} - f_i) = 0$  for  $i = 1, 2$ . Thus if  $\mu$  is not an eigenvalue of  $G_i$  then  $G_i$  is a clique or a co-clique because it has at most two distinct eigenvalues. In this case, by [10, Lemma 3]  $G_i$  is contained in a star complement for  $\mu$ , and so  $n_i \leq t$ . From the proof of Theorem 4.4 we see that for  $i, j \notin X_i$  there are just two values for  $|\Delta(i) \cap \Delta(j) \cap X_i|$  according as  $i \sim j$ ,  $i \not\sim j$ . Therefore  $n \leq \frac{1}{2}n_i(n_i - 1)$  by [5, Theorem 1.51]. Hence  $n_i = t$  and  $G_i$  is a star complement for  $\mu$  in  $G$ . If  $G_i$  is a clique then  $G = \Gamma_{36}$  by [8, Theorem 5.1].

Now suppose for definiteness that  $G_2$  is a co-clique. Then  $k_{22} = 0$  and  $k_{21} = d_2$ . From Eq.(2) we have  $\mu I - A_1 = \mu^{-1}B^\top B$ , whence  $k_{12} = \mu^2$ . Moreover, if  $i, j$  are adjacent vertices in  $X_1$  then  $|\Delta(i) \cap \Delta(j) \cap X_2| = -\mu$ ; and if  $i, j$  are non-adjacent vertices in  $X_1$  then  $|\Delta(i) \cap \Delta(j) \cap X_2| = 0$ . Since  $n_1 \neq t$ , the preceding argument shows that  $G_1$  is not a clique; and  $G_1$  is not a co-clique because  $G$  is not bipartite. It follows that  $|\Delta(i) \cap \Delta(j) \cap X_2|$  ( $i, j \in X_1$ ) takes precisely two values, and so by [5, Theorem 1.51] the neighbourhoods  $\Delta(i) \cap X_2$  ( $i \in X_1$ ) constitute a tight 4-design  $\mathcal{D}$  (see [5, p.20]). Note that  $\mu < 0$ , and so  $\mu^2 \geq 4$ . Also,  $\mu^2 \leq t - 2$  because the neighbourhoods  $\Delta(i) \cap X_2$  ( $i \in X_1$ ) are distinct. If  $\mu^2 < t - 2$  then by [5, Theorem 1.54]  $\mathcal{D}$  or its complement is the Steiner system  $S(4, 7, 23)$ . Therefore there are two cases to consider: (a)  $\mu = -4$  and  $t = 23$ , (b)  $\mu^2 = t - 2$ .

Now  $\rho\mu = \det K = -\mu^2 d_2$ , and so  $\rho = -\mu d_2$ . Since  $k_{22} = 0$ , Lemma 2.2 shows that  $a_1 d_2 = a_2 \rho$ , whence  $a_1 = -\mu a_2$  and  $-\mu a_2^2 = a_1 a_2 = -\mu(\lambda + 1)$ . Now we have

$$a_1^2 = \mu^2(\lambda + 1), \quad a_2^2 = \lambda + 1, \quad d_2 = \lambda + 1 - \lambda\mu, \quad \rho = -\lambda\mu - \mu + \lambda\mu^2.$$

By Lemma 2.2,

$$t = n_2 = \frac{(\rho + \lambda\mu - a_1^2)(\rho - \lambda)}{a_2^2 - a_1 a_2} = \frac{\mu(\mu d_2 + \lambda)}{\lambda + 1},$$

and so  $t - 1 = d_2(\mu^2 - 1)/(\lambda + 1)$ . On the other hand,  $\frac{1}{2}(t - 1) = n_1/n_2 = k_{21}/k_{12} = d_2/\mu^2$ .

In case (a) we have  $d_2 = \frac{1}{2}(t - 1)\mu^2 = 176$  and  $\rho = 704$ , while  $n = \frac{1}{2}t(t + 1) = 276$ , a contradiction. In case (b) we have  $2d_2 = (t - 1)\mu^2 = \mu^2 d_2(\mu^2 - 1)/(\lambda + 1)$ , whence  $\lambda = \frac{1}{2}(t - 1)(t - 4)$  and  $\rho = -\frac{1}{2}\mu(t - 1)(t - 2)$ . Now  $0 = \text{tr} A = \rho + \frac{1}{2}t(t - 1)\mu + (t - 1)\lambda$ , whence  $-2\mu = (t - 1)(t - 4)$  and  $4(t - 2) = 4\mu^2 = (t - 1)^2(t - 4)^2$ , a contradiction.  $\square$

It remains to consider the case in which  $\mu$  is an eigenvalue of both  $G_1$  and  $G_2$ . (A contradiction in this case would show that  $\mathcal{B} \cap \mathcal{C} = \{K_{1,2}, \Gamma_{36}\}$ .)

## 5 The remaining case

Here we assume that  $G$  is a graph in  $\mathcal{B} \cap \mathcal{C}$  other than  $\Gamma_{36}$ , so that  $\mu$  is an eigenvalue of both  $G_1$  and  $G_2$ . Moreover  $\mu \neq -2$  for otherwise  $\mu$  is the least



eigenvalue of  $G$  and then  $\Gamma_{36}$  is the only candidate for  $G$  (see [6, Chapter 6]). We take the distinct eigenvalues of  $G_i$  to be  $k_{ii}, \lambda_i$  and  $\mu$ . In this final section we offer four further observations which may be of interest to the reader who wishes to pursue matters further.

**Remark 5.1**  $|\mu + 1| \leq |\sqrt{d_1} - \sqrt{d_2}|$ .

**Proof.** Let  $\kappa = (\mu + 1)^2$ . By Proposition 4.2,  $\rho^2 = \rho(d_1 + d_2 - \kappa) - d_1 d_2$ , whence  $\kappa < d_1 + d_2$ . The quadratic  $\rho^2 - (d_1 + d_2 - \kappa)\rho + d_1 d_2$  has discriminant  $(d_1 + d_2 - \kappa)^2 - 4d_1 d_2$ , and so  $d_1 + d_2 - \kappa \geq 2\sqrt{d_1 d_2}$ , equivalently  $\kappa \leq (\sqrt{d_1} - \sqrt{d_2})^2$  as required.  $\square$

**Remark 5.2**  $\lambda_1 + \lambda_2 = \lambda + \mu$ .

**Proof.** We have  $\lambda_1 + \mu = e_1 - f_1$  and  $\lambda_2 + \mu = e_2 - f_2$ . By Theorem 4.4,  $(\lambda_1 + \lambda_2 + 2\mu)(d_2 - d_1) =$

$$\lambda(\mu^2 + \mu + \rho - d_1) + 2d_2\mu - d_1\mu + \rho - \lambda(\mu^2 + \mu + \rho - d_2) - 2d_1\mu - d_2\mu - \rho = (d_2 - d_1)(\lambda + 3\mu), \text{ and the result follows. } \square$$

**Remark 5.3** If  $\mathbf{x}$  is a  $\mu$ -eigenvector of  $G_1$  then  $\begin{pmatrix} \mathbf{x} \\ \mathbf{0} \end{pmatrix}$  is a  $\mu$ -eigenvector of  $G$ , and if  $\mathbf{y}$  is a  $\mu$ -eigenvector of  $G_2$  then  $\begin{pmatrix} \mathbf{0} \\ \mathbf{y} \end{pmatrix}$  is a  $\mu$ -eigenvector of  $G$ .

**Proof.** Let  $A = \begin{pmatrix} A_1 & M_1^\top \\ M_1 & A_2 \end{pmatrix}$ . From Eq.(1) we have

$$A_1^2 + M_1^\top M_1 - (\lambda + \mu)A_1 + \lambda\mu I = a_1^2 J.$$

Since  $\mathbf{j}^\top \mathbf{x} = 0$ , we have

$$\mu^2 \mathbf{x} + M_1^\top M_1 \mathbf{x} - (\lambda + \mu)\mu \mathbf{x} + \lambda\mu \mathbf{x} = \mathbf{0},$$

whence  $M_1 \mathbf{x} = \mathbf{0}$  and  $A \begin{pmatrix} \mathbf{x} \\ \mathbf{0} \end{pmatrix} = \mu \begin{pmatrix} \mathbf{x} \\ \mathbf{0} \end{pmatrix}$ . Similarly,  $A \begin{pmatrix} \mathbf{0} \\ \mathbf{y} \end{pmatrix} = \mu \begin{pmatrix} \mathbf{0} \\ \mathbf{y} \end{pmatrix}$ .  $\square$

**Remark 5.4** If  $d_1 < d_2$  and  $\mu < 0$  then  $d_2 - \rho < \frac{a_1 a_2 + \mu - \lambda}{a_2^2} \rho < \rho$ .

**Proof.** By Lemma 2.2 and Theorem 4.4 the inequality  $f_2 \leq k_{22}$  becomes

$$\frac{(\mu^2 + \mu + \rho - d_1)(d_2 + \lambda\mu)}{d_1 - d_2} \leq \frac{a_2 \rho - a_1 d_2}{a_2 - a_1},$$

equivalently

$$d_2 \mu + (\rho - d_1)\lambda + (\mu^2 + \mu)\lambda \geq -\rho.$$

Since  $\rho < d_2$ , we have  $(d_2 + \mu\lambda)(\mu + 1) + (\rho - d_1)\lambda > 0$ . Multiplying by  $\rho - d_2$ , and invoking Proposition 4.2, we obtain

$$d_2 - \rho < \frac{-(\mu + 1)\lambda\rho}{d_2 + \lambda\mu} = \frac{a_1 a_2 + \mu - \lambda}{a_2^2} \rho < \frac{a_1}{a_2} \rho < \rho.$$

$\square$

Remark 5.2 is useful in ruling out various candidates for prescribed  $t$  and  $\mu$ , as we illustrate below.

**Example 5.5** Here we show that  $(t, \mu) \neq (24, -3)$ . If  $t = 24$  then  $G_1, G_2$  are strongly regular graphs whose orders sum to 300. Taking  $n_1 \geq n_2$  without loss of generality, we find from Brouwer's list of feasible parameters [2] that when  $\mu = -3$  the only possibilities are:

- (i)  $(n_1, k_{11}, \lambda_1) = (175, 102, 17), (n_2, k_{22}, \lambda_2) = (125, 72, 12)$ ;
- (ii)  $(n_1, k_{11}, \lambda_1) = (231, 30, 9), (n_2, k_{22}, \lambda_2) = (69, 20, 5)$ ;
- (iii)  $(n_1, k_{11}, \lambda_1) = (275, 162, 27), (n_2, k_{22}, \lambda_2) = (25, 12, 2)$ .

Now  $\lambda = \lambda_1 + \lambda_2 + 3$  by Remark 5.2, and we can find  $\rho$  from the equation  $\rho + (t - 1)\lambda + \frac{1}{2}t(t - 1)\mu = 0$ . In each case we obtain the contradiction  $\rho + \mu \neq k_{11} + k_{22}$ . Analogous arguments eliminate the possibilities  $(t, \mu) = (24, -4), (23, -3), (23, -4)$ .  $\square$

**Acknowledgements** The author is grateful to Jack Koolen for his interest in the problem, and to the anonymous referees for their constructive comments.

## References

- [1] F. K. Bell and P. Rowlinson, On the multiplicities of graph eigenvalues, J. London Math. Soc. 35 (2003), 401-408.
- [2] A. E. Brouwer, A table of parameters of strongly regular graphs, <http://www.win.tue.nl/~aeb/>.
- [3] X-M. Cheng, A. L. Gavriluk, G. R. W. Greaves and J. H. Koolen, Biregular graphs with three eigenvalues, European J. Math. 56 (2016), 57-80.
- [4] X-M. Cheng, G. R. W. Greaves and J. H. Koolen, Graphs with three eigenvalues and second largest eigenvalue at most 1, J. Combinatorial Theory Ser. B 129 (2018), 55-78.
- [5] P. J. Cameron and J. H. van Lint, Designs, Graphs, Codes and their Links, Cambridge University Press (Cambridge), 1991.
- [6] D. Cvetković, P. Rowlinson and S. K. Simić, Spectral Generalizations of Line Graphs, Cambridge University Press (Cambridge), 2004.
- [7] D. Cvetković, P. Rowlinson and S. K. Simić, An Introduction to the Theory of Graph Spectra, Cambridge University Press (Cambridge), 2010.
- [8] D. Cvetković, P. Rowlinson and S. K. Simić, Graphs with least eigenvalue  $-2$ : ten years on, Linear Algebra Appl. 484 (2015), 504-539.

- [9] W. H. Haemers and D. G. Higman, Strongly regular graphs with strongly regular decomposition, *Linear Algebra Appl.* 114/115 (1989), 379-398.
- [10] M. Milošević. An example of using star complements in classifying strongly regular graphs, *Filomat* 22:2 (2008), 53-57.
- [11] P. Rowlinson, The main eigenvalues of a graph: a survey, *Applicable Analysis and Discrete Math.* 1 (2007), 445-471.
- [12] P. Rowlinson, On graphs with just three eigenvalues, *Linear Algebra Appl.* 507 (2016), 462-473.
- [13] P. Rowlinson, More on graphs with just three eigenvalues, *Applicable Analysis and Discrete Math.* 11 (2017), 74-80.